

(New) The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

In the Claims

Please amend the claims as follows. The claims are amended in accordance with 37 C.F.R. §1.121. No new matter has been added.

1. (Amended) A *in vitro* clonal propagating method for cultivation of marine algae, said method comprising the steps of:

a) establishing axenic viable material of algae for tissue culture by sequential treatment of the algal material in sterile sea water supplemented with domestic liquid detergent, povidine iodine and finally incubating the treated material in Provasoli enriched seawater (PES) medium with a broad spectrum antibiotic mixture and a fungicide for about 24 to 96 hours followed by thorough cleaning with sterile sea water to remove any traces of antibiotics and fungicide and blotting with sterile filter paper to obtain axenic explants;

b) culturing the axenic explants on agar plates fortified with PES medium at a temperature ranging between 20-25°C in the presence of cool white fluorescent lights at about 20-50 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$ irradiance and a 12 : 12 light dark cycle for induction of callus;

c) excising the callus from the explant after a period of at least 40 days and subculturing the callus on fresh agar plates fortified with PES medium in the presence of cool white fluorescent lights with 40-60 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$ irradiance and a 12 : 12 light and dark cycle to obtain differentiated densely pigmented oval or spherical shaped micro-propagules;

d) subculturing thin slices of the pigmented callus as an embedded culture in agar plates in Provasoli Enriched Seawater (PES) medium containing plant growth regulators, for a period of about 20 to 40 days, in the presence of cool white fluorescent lights of 20-60 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$ irradiance and a 12 : 12 light and dark cycle to achieve profusely branched pigmented calli in each embedded block leading to enhanced somatic embryogenesis and micro-propagule formation in pigmented filamentous callus;

e) transferring the filamentous calli with somatic embryos to liquid PES medium in an agitated condition for morphogenesis and development of young plantlets with multiple shoots from propagules; and

f) cultivating algal biomass on a large scale in the sea by growing the young plantlets in enclosed perforated polythene bags.

2. (Amended) A method as claimed in claim 1, wherein the material for tissue culture is a Rhodophytic marine alga selected from the group of genera of *Eucheuma*, *Gigartina*, and *Chondrus*.

3. (Amended) A method as claimed in claim 1, wherein the material for tissue culture is an alga selected from the group of *Eucheuma striatum*, *Kappaphycus alvarezii*, *Eucheuma cottonii*, *Eucheuma denticulatum*, *Eucheuma spinosum*, *Eucheuma alvarezii*, *Eucheuma procrusteanum*, *Gigartina intermedia*, *Gigartina exasparata* and *Chondrus crispus*.

8. (Amended) A method as claimed in claim 1, wherein the calli are subcultured by growing thin slices of pigmented calli as embedded cultures in agar plates containing 0.3-0.6% agar and made in provasoli enriched seawater medium at 20-25°C in the presence of cool white fluorescent light at about 20-50 μ mol photon $m^{-2} s^{-1}$ irradiance with 12:12 light and dark cycle to obtain profusely branched filamentous pigmented calli in each embedded block.

15. (Amended) A method as claimed in claim 1, wherein said step of subculturing thin slices of the pigmented callus includes adding growth regulators including α -naphthalene acetic acid and 6-benzylaminopurine to achieve further enhancement of formation of somatic embryos through somatic embryogenesis.

16. (Amended) A method as claimed in claim 1, wherein a harvesting period after at least 60 days can yield a higher biomass of tissue cultured plant than that of a control of parent plants or wherein the biomass can be maintained constant and a cultivation period reduced from at least 60 days.

17. (Amended) A method as claimed in claim 1, wherein a two fold increase in growth in fresh weight of tissue cultured plant is achieved over a control of parent plants, without change in carrageenan product yield and gel strength, through micro-propagule formation from pigmented calli.

18. (Amended) A method as claimed in claim 1, wherein the material for tissue culture is a Phaeophytic marine alga selected from the group of genera of *Luminaria*, *Undaria*, *Ecklonia*, *Eisenia*, *Macrocystis*, *Sargassum*, and *Turbinaria*.